

Dynamics of Weyl Scale Invariant non-BPS $p=3$ Brane

Lu-Xin Liu

*Department of Physics, Purdue University,
West Lafayette, IN 47907, USA*

and

*National Institute for Theoretical Physics,
Department of Physics and Centre for Theoretical Physics,
University of the Witwatersrand,
Wits, 2050, South Africa
Luxin.Liu9@gmail.com*

Abstract. The action of Weyl scale invariant $p=3$ brane which breaks the target super Weyl scale symmetry in the $N=1$, 5D superspace down to the lower dimensional Weyl symmetry $W(1,3)$ is derived by the approach of coset method. The infeasibility of embedding an alternative Minkowski brane in the target super Weyl spacetime is also pointed out.

In Type II string theory, BPS D-branes which carry R-R charges break the bulk $N=2$ spacetime supersymmetries and are invariant under only half of the original supersymmetries [1]. Its supersymmetric extension of Born-Infeld action describing the super BPS D-branes is realized by using Kappa-symmetry, which is needed for gauging away one half of the fermionic degrees of freedom [2]. In addition, super BPS D-branes theories in dimensions other than ten were discussed in Ref.[3]. However, when no Kappa-symmetry is considered, the number of fermionic degree of freedom becomes the appropriate number of fermionic Goldstone modes for completely breaking the bulk supersymmetry. This is the case of non-BPS D-brane[4]. Although there is no manifest supersymmetry for this non-BPS D-brane world volume theory, the world volume theory is supersymmetric but realized as a spontaneously broken symmetry.

There have been many generalized proposals for the construction of non-BPS D-branes [5]. In Ref.[6], a space filling non-BPS D-brane, which breaks the supersymmetries of the target non-centrally extended $N=2$, 3D superspace, is introduced by means of the Green-Schwarz approach [7]. This action, which is found to be dual to a non-BPS $p=2$ brane, can be alternatively constructed from coset approach and totally breaks the supersymmetry in the embedded $N=1$, 4D superspace.

In this paper, we consider a $p=3$ brane embedded in 5D target bulk space with super-Weyl symmetry, which includes $N=1$, 5D super-Poincare symmetry and the Weyl scale symmetry. Since there is no observation for the superpartners for all the particles in the standard model of particle physics, the supersymmetry must be a broken one for a realistic theory. On the other hand, in particle physics it has been well known that the scale symmetry plays an important role [8-12], and it has been applied to a wide range of physics sectors, which include string theory, branes [13-16], as well as recent unparticle physics theory [17]. It is notable that the scale invariance is manifestly broken by the masses of the particles in the standard model; nevertheless, it is conceivable that at a much higher energy scale beyond the standard model, there are nontrivial sectors in particle physics that have the property of scale invariance. Accordingly, we consider that the dynamics of the embedded $p=3$ brane is described by Weyl scale theory, which also totally breaks the supersymmetry in the target bulk space.

The dynamics of the brane describes the long wave oscillation modes of the brane into the target bulk space. In this regards, we use the coset approach to realize the sponetaneous breaking of the target bulk symmetries down to the subgroup symmetries on the brane world volume. Consider the 5D dimension super-Weyl group G . Its generators include $W(1, 4)$ [8] Weyl group (formed by M^{MN} , P^M and Weyl scale (dilatation) generator D) and eight minimum supersymmetrical charges Q_a with four complex components. The component index $a=1, 2, 3, 4$. Its algebra has the following (anti)commutation relations:

$$\begin{aligned}
\{Q_a, \bar{Q}_b\} &= 2\gamma_{ab}^M P_M \\
[Q_a, M^{MN}] &= \frac{1}{2}\gamma_{ab}^{MN} Q_b \\
[M_{MN}, M_{OQ}] &= i(\eta_{NO}M_{MQ} + \eta_{MQ}M_{NO} - \eta_{MO}M_{NQ} - \eta_{NQ}M_{MO}) \\
[M_{MN}, P_R] &= i(\eta_{NR}P_M - \eta_{MR}P_N)
\end{aligned} \tag{1}$$

and

$$\begin{aligned}
[D, M_{MN}] &= 0, [D, D] = 0, [D, P_M] = -iP_M \\
[D, Q_a] &= -\frac{1}{2}iQ_a, [D, \bar{Q}_a] = -\frac{1}{2}i\bar{Q}_a,
\end{aligned} \tag{2}$$

where $\eta_{MN} = (1, -1, -1, -1, -1)$, $\gamma^M = (\gamma^\mu, \gamma^4)$, $\gamma^4 = -\gamma^0\gamma^1\gamma^2\gamma^3$, $\bar{Q}_a = Q_b^+ \gamma_{ba}^0$, $\gamma^{MN} = i[\gamma^M, \gamma^N]/2$ and $M, N=0, 1, 2, 3, 4$. In the present context, we consider a $p=3$ brane embedded in the $N=1$, 5D superspace. Accordingly, the three spacial dimensional brane breaks down the target space super-Weyl invariance to a lower dimensional Weyl group symmetry $W(1, 3)$, whose unbroken generators are $\{M_{\mu\nu}, D, P_\mu\}$, where the index $\mu, \nu = 0, 1, 2, 3$. As a result, the broken generators, besides the spontaneously broken automorphism generators $M_{\mu 4}$, include the translational generator P_4 transverse to the brane directions and the spinorial generators Q_a and $\bar{Q}_a$ related to the Grassmann coordinate directions in the superspace. On the other hand, one may be tempted to consider the Lorentz group as the stability group for the total symmetry breaking. However, at the end of the paper, we point out that infeasibility of embedding such a Minkowski brane in this super-Weyl spacetime. Since there would be the dilaton field localized on the brane world volume corresponding to the target scale symmetry breaking, and one will find it contributes a potential term on the brane world volume with an unbounded VEV. This fact then excludes

the brane modes that simultaneously nonlinear realize dilatation symmetry and the supersymmetry.

We work on the 4D world volume of the submanifold. The 5D super Weyl algebra is to be converted to 4D algebra through dimension reduction. In the Weyl representation, we introduce two supersymmetrical Weyl spinor charges Q_α and S_α and define

$$Q_a = \frac{1}{\sqrt{2}} \begin{pmatrix} Q_\alpha \\ i\bar{S}^{\dot{\alpha}} \end{pmatrix}, \bar{Q}_a = (-iS^\alpha, \bar{Q}_{\dot{\alpha}}) \quad (3)$$

where the Weyl spinor indices $\alpha, \dot{\alpha} = 1, 2$. Therefore, the algebra becomes

$$\begin{aligned} \{Q, \bar{Q}\} &= 2\sigma^\mu P_\mu, \{S, \bar{S}\} = 2\sigma^\mu P_\mu, \\ \{Q, S\} &= -2\varepsilon P_4 = -2\varepsilon Z, \\ [K^\mu, Q] &= -i\sigma^\mu \bar{S}, [K^\mu, S] = i\sigma^\mu \bar{Q}, \\ [M^{\mu\nu}, Q] &= -\frac{1}{2}\sigma^{\mu\nu} Q, [M^{\mu\nu}, S] = -\frac{1}{2}\sigma^{\mu\nu} S, \end{aligned} \quad (4)$$

along with their Weyl scale properties

$$\begin{aligned} [D, K_\mu] &= 0, [D, Z] = -iZ, [D, P_\mu] = -iP_\mu \\ [D, Q_\alpha] &= -\frac{1}{2}iQ_\alpha, [D, S_\alpha] = -\frac{1}{2}iS_\alpha \end{aligned} \quad (5)$$

where $K^\mu = 2M^{4\mu}$. Therefore, from the 4D standpoint of view, the N=1, 5D SUSY algebra is a central-charged N=2 four dimensional extended superalgebra with one 5D translation generator becoming the central charge.

Here, the target symmetry group G is restricted to the group whose generators can be divided into two subgroups with one is the automorphism group of another. The transformation of the group G with respect to the coset of the unbroken automorphism generator group would give us a description of the vierbein of the embedded submanifold, which has the same dimension as the coset space with respect to the unbroken automorphism group of the unbroken whole subgroup. In our case, the coset is G/H , where H is formed by unbroken automorphism generators $\{M_{\mu\nu}, D\}$. Hence the embedded three spatial dimension brane has $W(1, 3)$ symmetry structure oscillating through the coset space formed by G/H' , where H' is spanned by the automorphism generators $\{M_{MN}, D\}$ and is the automorphism group of the super

spacetime group formed by the set $\{Q_\alpha, \bar{Q}_{\dot{\alpha}}, S_\alpha, \bar{S}_{\dot{\alpha}}, P_\mu\}$. The submanifold which the brane sweeps out has the dimensions of the coset G'/H with tangent space group H , where G' is spanned by the unbroken automorphism generators $\{M_{\mu\nu}, D\}$ and the unbroken spacetime translation generators P_μ . The group elements of G can be parameterized in some neighborhood of the identity element as follows

$$g = e^{i[a^\mu p_\mu + \xi Q + \bar{\xi} \bar{Q} + \psi S + \bar{\psi} \bar{S} + z Z + b^\mu K_\mu + \alpha^{\mu\nu} M_{\mu\nu} + dD]} \quad (6)$$

We choose static gauge for the parameterization of the brane world volume, then the space time coordinates x^μ lie in the brane parameterizing direction ξ^μ . The coset G/H representative elements then have the form

$$\Omega = e^{ix^\mu p_\mu} e^{i\phi(x)Z} e^{i[\theta(x)Q + \bar{\theta}(x)\bar{Q} + \lambda(x)S + \bar{\lambda}(x)\bar{S}]} e^{iu^\mu(x)K_\mu} \quad (7)$$

in which $\phi(x), \theta(x), \bar{\theta}(x), \lambda(x), \bar{\lambda}(x)$ and $u(x)$ are the Nambu-Goldstone fields corresponding to each broken generators. Since the elements of group G can be decomposed uniquely into a product form, given by a coset representative element and a subgroup element h , the transformations of these Nambu-Goldstone fields can be obtained by acting a group operation on Ω from the left, i.e.

$$g\Omega = \Omega' h \quad (8)$$

where the new coset element $\Omega' = e^{ix'^\mu p_\mu} e^{i\phi'(x')Z} e^{i[\theta'(x')Q + \bar{\theta}'(x')\bar{Q} + \lambda'(x')S + \bar{\lambda}'(x')\bar{S}]} e^{iu'^\mu(x')K_\mu}$.

As for the spontaneous symmetry breaking of (super)spacetime, the Nambu-Goldstone fields are those associated with the broken (super)translation operators, and the superfluous Nambu-Goldstone field u^μ can be eliminated by imposing covariant constraints on the Cartan differential one forms. Explicitly, the Cartan one-forms can be expanded with respect to the full generators as follows

$$\begin{aligned} \Omega^{-1} d\Omega &= e^{-iu^\mu(x)K_\mu} e^{-i[\theta(x)Q + \bar{\theta}(x)\bar{Q} + \lambda(x)S + \bar{\lambda}(x)\bar{S}]} e^{-i\phi(x)Z} e^{-ix^\mu p_\mu} \\ &\quad d(e^{ix^\mu p_\mu} e^{i\phi(x)Z} e^{i[\theta(x)Q + \bar{\theta}(x)\bar{Q} + \lambda(x)S + \bar{\lambda}(x)\bar{S}]} e^{iu^\mu(x)K_\mu}) \\ &= i(\omega^a p_a + \omega_Q^\alpha Q_\alpha + \bar{\omega}_{\bar{Q}\dot{\alpha}} \bar{Q}^{\dot{\alpha}} + \omega_Z Z + \omega_S^\alpha S_\alpha + \bar{\omega}_{\bar{S}\dot{\alpha}} \bar{S}^{\dot{\alpha}} + \\ &\quad \omega_k^a K_a + \omega_M^{\mu\nu} M_{\mu\nu} + \omega_D D) \end{aligned} \quad (9)$$

According to Eq.(8), the Cartan one-forms transform as

$$\Omega'^{-1} d\Omega' = h(\Omega^{-1} d\Omega)h^{-1} + h dh^{-1} \quad (10)$$

By using the differentiation formula for exponent $\exp(-b)d\exp(b) = \sum_{k=0}^{\infty} \frac{(-1)^k}{(k+1)!} (ad_b)^k db$, where $ad_b(a) = [b, a]$ is the adjoint operation, we get the following building blocks related to the construction of the effective brane action, which is invariant under G transformations:

$$\begin{aligned}
\omega^a &= (dx^b + i\theta\sigma^b d\bar{\theta} - id\theta\sigma^b \bar{\theta} + i\lambda\sigma^b d\bar{\lambda} - id\lambda\sigma^b \bar{\lambda}) \\
&\quad \cdot (\delta_b^a + (\cosh 2\sqrt{u^2} - 1) \frac{u_b u^a}{u^2}) - (d\phi + id(\theta\lambda - \bar{\theta}\bar{\lambda})) \frac{\sinh 2\sqrt{u^2}}{\sqrt{u^2}} u^a \\
\omega_Z &= (d\phi + id(\theta\lambda - \bar{\theta}\bar{\lambda})) \cosh 2\sqrt{u^2} \\
&\quad - (dx^a + i(\theta\sigma^a d\bar{\theta} - d\theta\sigma^a \bar{\theta} + \lambda\sigma^a d\bar{\lambda} - d\lambda\sigma^a \bar{\lambda})) \frac{\sinh 2\sqrt{u^2}}{\sqrt{u^2}} u_a \\
\omega_D &= 0
\end{aligned} \tag{11}$$

where $a = 0, 1, 2, 3$. We use Latin letters a, b to represent the tangent spacetime indices, and Greek letters μ, ν to denote 1 + 3 general coordinate indices in what follows. The vierbein can be found by expanding the Cartan one-forms associated with the unbroken spacetime generators with respect to the general coordinate differentials dx^μ , i.e. $\omega^a = dx^\mu e_\mu^a$. The spin connection $\omega_{\rho M}^{\mu\nu}$ can also be constructed in a similar way, i.e., $\omega_M^{\mu\nu} = dx^\rho \omega_{\rho M}^{\mu\nu}$. Considering the unbroken subgroup $W(1, 3)$ (Weyl group) and Eq.(10), it follows that the dilatation transformation property of the vierbein takes the form

$$\omega^a \rightarrow e^d \omega^a \text{ or } e_\mu^a \rightarrow e^d e_\mu^a \tag{12}$$

then it is concluded that it has scale dimension 1. On the brane world volume, the interval is

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu = \eta_{ab} e_\mu^a e_\nu^b dx^\mu dx^\nu \tag{13}$$

in which the metric tensor is given by

$$g_{\mu\nu} = e_\mu^a e_\nu^b \eta_{ab}. \tag{14}$$

Besides, under the scale transformation, in the local tangent space we have the transformation

$$x^a \rightarrow x'^a = e^d x^a \tag{15}$$

it follows that the interval transforms as

$$ds^2 \rightarrow ds'^2 = e^{2d} ds^2 \quad (16)$$

This can also be concluded from Eqs.(12) and (13). In terms of metric tensor $g_{\mu\nu}$, considering Eq.(12), it is understanding that the metric tensor is weight 2 under the Weyl scale transformation, i.e.

$$g'_{\mu\nu} = e^{2d} g_{\mu\nu}. \quad (17)$$

Conversely, $g^{\mu\nu}$ has scale weight -2 . Explicitly, in accord with Eq.(11), the vierbein has the form

$$\begin{aligned} e_\mu^a &= (\delta_\mu^b + i\theta\sigma^b\partial_\mu\bar{\theta} - i\partial_\mu\theta\sigma^b\bar{\theta} + i\lambda\sigma^b\partial_\mu\bar{\lambda} - i\partial_\mu\lambda\sigma^b\bar{\lambda}) \\ &\quad \cdot (\delta_b^a + (\cosh 2\sqrt{u^2} - 1)\frac{u_b u^a}{u^2}) - \partial_\mu(\phi + i\theta\lambda - i\bar{\theta}\bar{\lambda})\frac{\sinh 2\sqrt{u^2}}{\sqrt{u^2}}u^a \\ &= A_\mu^b \cdot (\delta_b^a + (\cosh 2\sqrt{u^2} - 1)\frac{u_b u^a}{u^2} - \tilde{D}_b(\phi + i\theta\lambda - i\bar{\theta}\bar{\lambda})\frac{\sinh 2\sqrt{u^2}}{\sqrt{u^2}}u^a) \end{aligned} \quad (18)$$

in which $A_\mu^b = (\delta_\mu^b + i\theta\sigma^b\partial_\mu\bar{\theta} - i\partial_\mu\theta\sigma^b\bar{\theta} + i\lambda\sigma^b\partial_\mu\bar{\lambda} - i\partial_\mu\lambda\sigma^b\bar{\lambda})$, and $\tilde{D}_b = A_b^{-1\mu}\partial_\mu$ is the Akulov-Volkov derivative[18-21]. Imposing the covariant condition $\omega_Z = 0$ on the Cartan one-forms, as a result of the inverse Higgs Mechanism [22], the field u_m can be eliminated by the following relation

$$u_b \frac{\tanh 2\sqrt{u^2}}{\sqrt{u^2}} = \tilde{D}_b\phi + i\tilde{D}_b\theta\lambda + i\theta\tilde{D}_b\bar{\lambda} - i\tilde{D}_b\bar{\theta}\bar{\lambda} - i\bar{\theta}\tilde{D}_b\bar{\lambda} = \tilde{D}_b\Phi \quad (19)$$

where $\Phi = \phi + i\theta\lambda - i\bar{\theta}\bar{\lambda}$. It can further yield

$$\cosh 2\sqrt{u^2} = \sqrt{\frac{1}{1 - \tanh^2 2\sqrt{u^2}}} = \sqrt{\frac{1}{1 - (\tilde{D}\Phi)^2}} \quad (20)$$

with

$$\tanh^2 2\sqrt{u^2} = (\tilde{D}\Phi)^2. \quad (21)$$

Plugging Eq.(19) back into Eq.(18), the vierbein hence has the simple form

$$\begin{aligned} e_\mu^a &= A_\mu^b \cdot (\delta_b^a + \frac{1 - \cosh 2\sqrt{u^2}}{\cosh 2\sqrt{u^2}} \frac{u_b u^a}{u^2}) \\ &= A_\mu^b \cdot (\delta_b^a + \frac{1 - \cosh 2\sqrt{u^2}}{\cosh 2\sqrt{u^2}} \frac{\tilde{D}_b\Phi\tilde{D}^a\Phi}{\tanh^2 2\sqrt{u^2}}) \\ &= A_\mu^b \cdot (\delta_b^a + \frac{\tilde{D}_b\Phi\tilde{D}^a\Phi}{(\tilde{D}\Phi)^2} (\sqrt{1 - (\tilde{D}\Phi)^2} - 1)). \end{aligned} \quad (22)$$

The metric tensor hence becomes

$$g_{\mu\nu} = e_\mu^a e_\nu^b \eta_{ab} = A_\mu^a A_\nu^b \eta_{ab} - \partial_\mu \Phi \partial_\nu \Phi. \quad (23)$$

Introduce five dynamic variables $X^M = (X^a, X^4) = (X^a, \Phi)$, which are defined as follows

$$\begin{aligned} dX^a &= dx^\mu A_\mu^a; \\ dX^4 &= \partial_\mu (\phi + i\theta\lambda - i\bar{\theta}\bar{\lambda}) dx^\mu. \end{aligned} \quad (24)$$

Therefore, in the static gauge $\xi^\mu = x^\mu$, the metric tensor in Eq.(23) gives us

$$\begin{aligned} g_{\mu\nu} &= \eta_{MN} \frac{\partial X^M}{\partial \xi^\mu} \frac{\partial X^N}{\partial \xi^\nu} \\ &= \eta_{\mu\nu} + i\theta\sigma_\nu \overset{\leftrightarrow}{\partial}_\mu \bar{\theta} + i\lambda\sigma_\nu \overset{\leftrightarrow}{\partial}_\mu \bar{\lambda} + i\theta\sigma_\mu \overset{\leftrightarrow}{\partial}_\nu \bar{\theta} + i\lambda\sigma_\mu \overset{\leftrightarrow}{\partial}_\nu \bar{\lambda} \\ &\quad + (i\theta\sigma_b \overset{\leftrightarrow}{\partial}_\mu \bar{\theta} + i\lambda\sigma_b \overset{\leftrightarrow}{\partial}_\mu \bar{\lambda}) \cdot (i\theta\sigma^b \overset{\leftrightarrow}{\partial}_\nu \bar{\theta} + i\lambda\sigma^b \overset{\leftrightarrow}{\partial}_\nu \bar{\lambda}) \\ &\quad - (\partial_\mu \phi + i\partial_\mu \theta\lambda + i\theta\partial_\mu \lambda - i\partial_\mu \bar{\theta}\bar{\lambda} - i\bar{\theta}\partial_\mu \bar{\lambda}) \\ &\quad \cdot (\partial_\nu \phi + i\partial_\nu \theta\lambda + i\theta\partial_\nu \lambda - i\partial_\nu \bar{\theta}\bar{\lambda} - i\bar{\theta}\partial_\nu \bar{\lambda}) \\ &= \eta_{MN} \frac{\partial x^M}{\partial \xi^\mu} \frac{\partial x^N}{\partial \xi^\nu} + \theta, \bar{\theta}, \lambda, \text{ and } \bar{\lambda} \text{ terms} \end{aligned} \quad (25)$$

where $x^M = (x^0, x^1, x^2, x^3, \phi)$. Consequently, in contrast with the normal induced metric $g_{\mu\nu} = \eta_{MN} \frac{\partial x^M}{\partial \xi^\mu} \frac{\partial x^N}{\partial \xi^\nu}$ on the p-brane world volume, there are modification terms to the metric, which are contributed by the Nambu-Goldstone fields $\theta(x), \bar{\theta}(x), \lambda(x)$, and $\bar{\lambda}(x)$, as a result of the broken symmetries associated with the superspace coordinate directions.

Introduce an auxiliary (intrinsic) metric $\rho_{\mu\nu}$ on the p=3 brane world volume, whose scale dimension is 2 as induced by the scale transformation $\xi^\mu \rightarrow e^d \xi^\mu$, i.e.

$$\rho_{\mu\nu} \rightarrow e^{2d} \rho_{\mu\nu} \quad (26)$$

or

$$dx^4 \sqrt{|\rho|} \rightarrow dx'^4 \sqrt{|\rho'|} = e^{4d} dx^4 \sqrt{|\rho|}. \quad (27)$$

By using Eq.(25), the effective scale invariant action of the p=3 brane world volume

is secured as follows

$$\begin{aligned}
I &= -T \int d^4x \sqrt{|\rho|} \left[\frac{1}{4} \rho^{\mu\nu} \eta_{MN} \partial_\mu X^M \partial_\nu X^N \right]^2 \\
&= -T \int d^4x \sqrt{-\rho} \left[\frac{1}{4} \rho^{\mu\nu} (\eta_{\mu\nu} + i\theta \sigma_\nu \overset{\leftrightarrow}{\partial}_\mu \bar{\theta} + i\lambda \sigma_\nu \overset{\leftrightarrow}{\partial}_\mu \bar{\lambda} + i\theta \sigma_\mu \overset{\leftrightarrow}{\partial}_\nu \bar{\theta} + i\lambda \sigma_\mu \overset{\leftrightarrow}{\partial}_\nu \bar{\lambda} \right. \\
&\quad + (i\theta \sigma_b \overset{\leftrightarrow}{\partial}_\mu \bar{\theta} + i\lambda \sigma_b \overset{\leftrightarrow}{\partial}_\mu \bar{\lambda}) \cdot (i\theta \sigma^b \overset{\leftrightarrow}{\partial}_\nu \bar{\theta} + i\lambda \sigma^b \overset{\leftrightarrow}{\partial}_\nu \bar{\lambda}) \\
&\quad - (\partial_\mu \phi + i\partial_\mu \theta \lambda + i\theta \partial_\mu \lambda - i\partial_\mu \bar{\theta} \bar{\lambda} - i\bar{\theta} \partial_\mu \bar{\lambda}) \\
&\quad \cdot (\partial_\nu \phi + i\partial_\nu \theta \lambda + i\theta \partial_\nu \lambda - i\partial_\nu \bar{\theta} \bar{\lambda} - i\bar{\theta} \partial_\nu \bar{\lambda}) \Big]^2
\end{aligned} \tag{28}$$

in which the auxiliary intrinsic metric $\rho_{\mu\nu}$ can be eliminated by using its equation of motion. Here, $\rho^{\mu\nu}$ is the inverse of the metric $\rho_{\mu\nu}$ with scale dimension -2 , and ρ stands for the determinant of $\rho_{\mu\nu}$, and T stands for the brane tension. The part inside the square brackets has a scale dimension -2 . It can be concluded that Eq.(28) is Weyl scale invariant under the transformation $\xi^\mu \rightarrow e^d \xi^\mu$. Obviously, when the spinors are set to zero, it reduces to the Weyl scale invariant bosonic action [15]:

$$I_B = -\frac{f_s^2}{2} \int d^4\xi \sqrt{|\det G|} \left[\frac{1}{4} G^{\mu\nu} \eta_{ab} \frac{\partial x^a}{\partial \xi^\mu} \frac{\partial x^b}{\partial \xi^\nu} \right]^2 \tag{29}$$

The action (28) describes the effective oscillation modes of the brane into the bulk space, corresponding to the symmetry breakings in the $\theta, \bar{\theta}, \lambda, \bar{\lambda}, \phi$ (super)space coordinates directions, and whose long wave length excitation modes are described by these Nambu-Goldstone fields associated with those broken symmetries, i.e., the symmetries related to $Q_\alpha, \bar{Q}_{\dot{\alpha}}, S_\alpha, \bar{S}_{\dot{\alpha}}$, and Z .

On the other hand, if the embedded p=3 brane breaks the target scale symmetry as well, i.e., the coset representative elements have the following alternative form

$$\Omega_s = e^{ix^\mu p_\mu} e^{i\phi(x)Z} e^{i[\theta(x)Q + \bar{\theta}(x)\bar{Q} + \lambda(x)S + \bar{\lambda}(x)\bar{S}]} e^{iu^\mu(x)K_\mu} e^{i\sigma(x)D}. \tag{30}$$

Hence, the Cartan one-forms

$$\begin{aligned}
\Omega_s^{-1} d\Omega_s &= i(\omega_s^a p_a + \omega_{sQ}^\alpha Q_\alpha + \bar{\omega}_{s\bar{Q}\dot{\alpha}} \bar{Q}^{\dot{\alpha}} + \omega_{sZ} Z + \omega_{sS}^\alpha S_\alpha + \\
&\quad \bar{\omega}_{s\bar{S}\dot{\alpha}} \bar{S}^{\dot{\alpha}} + \omega_{sk}^a K_a + \omega_{sM}^{\mu\nu} M_{\mu\nu} + \omega_{sD} D)
\end{aligned} \tag{31}$$

gives us

$$\begin{aligned}
\omega_s^a &= \omega^a e^{-\sigma} = e^{-\sigma} (dx^b + i\theta\sigma^b d\bar{\theta} - id\theta\sigma^b \bar{\theta} + i\lambda\sigma^b d\bar{\lambda} - id\lambda\sigma^b \bar{\lambda}) \\
&\cdot (\delta_b^a + (\cosh 2\sqrt{u^2} - 1) \frac{u_b u^a}{u^2}) - (d\phi + id(\theta\lambda - \bar{\theta}\bar{\lambda}) \frac{\sinh 2\sqrt{u^2}}{\sqrt{u^2}} u^a) \\
&= dx^\mu e_{s\mu}^a \\
\omega_{sZ} &= \omega_Z e^{-\sigma} \\
\omega_{sD} &= dx^\mu \partial_\mu \sigma = \omega_s^a e_{sa}^{-1\mu} \partial_\mu \sigma = \omega_s^a D_\mu \sigma
\end{aligned} \tag{32}$$

in which, the covariant derivative $D_\mu = e_{sa}^{-1\mu} \partial_\mu$. Further, considering Eq.(8), for a pure scale transformation, we find the dilaton field transforms as

$$\sigma \rightarrow \sigma' = \sigma + d \tag{33}$$

Accordingly, the dilaton field, which transforms with a shift, behaves as the Nambu-Goldstone field, signaling the sponetaneous breaking of the scale symmetry in the target bulk space. Therefore, considering Eq.(32), the effective brane action, which includes all the Nambu-Goldstone modes, is found to be

$$\begin{aligned}
I &= -T \int d^4x \det e_s - \frac{T_s}{2} \int d^4x \det e_s \eta^{ab} D_a \sigma D_b \sigma \\
&= -T \int d^4x e^{-4\sigma} \det e - \frac{T_s}{2} \int d^4x \det e e^{-2\sigma} \eta^{ab} e_a^{-1\mu} e_b^{-1\nu} \partial_\mu \sigma \partial_\nu \sigma
\end{aligned} \tag{34}$$

in which T_s is related to the broken scale of the dilatation symmetry. Expanding the first term explicitly, we then have the potential

$$V \propto e^{-4\sigma} \tag{35}$$

Therefore, the VEV can be determined by estimating the value of $\langle e^{-4\sigma} \rangle$, which becomes minimum when $\langle \sigma \rangle$ goes to ∞ . However, due to the unbound of the VEV of the dilaton field it follows that σ can not be a NG particle. This fact thus indicates that it is infeasible to embed such a Minkowski brane, whose modes break the bulk dilatation symmetry and the supersymmetry simultaneously, in the target Super-Weyl bulk space.

In summary, in this paper, we start with the super Weyl group and its automorphism subgroup. Then the embedded Weyl scale invariant p=3 brane breaks the

super Weyl group symmetry in the target bulk space down to the $1+3$ dimensional Weyl $W(1,3)$ symmetry in the submanifold. The brane's dynamics is described by the Goldstone bosons (Goldstino fermions) modes associated with the broken spatial(Grassmann) generators of the symmetry(supersymmetry) group. The approach of nonlinear realization has been extensively used to describe the spontaneous partial breaking of (extended) supersymmetry and construct actions of (super)brane dynamics[23-27]. It also has been applied to branes of M theory with a large automorphism group of superalgebra [28]. In addition, in the theory of brane-world scenarios, the universe can be regarded as a four dimensional topological defect in the form of domain wall embedded in a higher dimensional spacetime [29], and it has been shown that the non-BPS topological defects can be a source of possible SUSY breaking [30]. Actually, it is found that the $N=1$ supersymmetry preserved on the four dimensional world volume of one wall (brane) is completely broken by the coexistence of the other wall (brane)[31]. As a result, it admits another possible origin of SUSY breaking, except those fluctuation modes of the brane in the super-space as discussed here.

The author thanks the THEP groups at Purdue and NITheP for supports.

REFERENCES

- [1] J. Polchinski, Phys. Rev. Lett. 75, 4724 (1995)
- [2] M.Aganagic, C.Popescu, and J.H.Schwarz, Phys.Lett.B393, 311(1997); M.Cederwall, A. von Gussich, B.E.W.Nilsson and A.Westerberg, Nucl.Phys.B490, 163(1997); E.Bergshoeff and P.K.Townsend, Nucl.Phys.B490, 145(1997); M.Gederwall, A.von Gussich, B.E.W.Nilsson, P.Sundellm, and A.Westerberg, Nucl.Phys.B490,179(1997); Igor A. Bandos and D.P.Sorokin, Nucl.Phys.B497, 275(1997)
- [3] E.Ivanov and S.Krivonos, Phys.Lett.B453, 237(1999); Sergei V. Ketov, Mod.Phys. Lett.A14, 501(1999); E.Ivanov; AIP Conf. Proc. 589, 61(2001)
- [4] A.Sen, JHEP 10, 008(1999)
- [5] See Review, for example, Matthias R. Gaberdiel, Class.Quant.Grav.17,3483(2000); A.Sen, hep-th/9904207
- [6] T. E. Clark, M. Nitta and T. ter Veldhuis, Phys.Rev.D70, 105005(2004)
- [7] M.B.Green and J.H.Schwarz, Phys.Lett. B136, 367(1984); M.B.Green, J.H.Schwarz and E.Witten, Superstring Theory (Cambridge Univ. Press, Cambridge, England, 1987) Vol.2
- [8] M. Blagojevic, Gravitation and Gauge Symmetries, IOP Publishing, 2002
- [9] Ryoyu Utiyama, Progress of Theor. Phys.50, 2080, 1973; Progress of Theor. Phys.53, 565, 1975
- [10] Kenji Hayashi and T.Kugo, Progress of Theor. Phys.61, 334, 1979
- [11] A. Bregman, Progress of Theor. Phys.49, 667, 1973
- [12] Kenji Hayashi, M.Kasuya and T.Shirafuji, Progress of Theor. Phys.57, 431, 1977
- [13] A.Aurilia, A.Smailagic and E.Spallucci, Phys.Rev.D51:4410, 1995
- [14] J.Antonio Garcia, Roman Linares and J.David Vergara, Phys. Lett. B503, 154(2001)

- [15] C.Alvear, R.Amorim and J.Barcelos-Neto, Phys. Lett. B273, 415(1991)
- [16] Lu-Xin Liu, Phys. Rev. D74, 45030(2006)[arXiv:hep-th/0602180]
- [17] H. Georgi, Phys.Rev.Lett.98:221601, 2007; Phys.Lett.B650:275-278, 2007
- [18] Lu-Xin Liu, Mod. Phys. Lett. A20, 2545(2005)[arXiv:hep-ph/0408210]
- [19] T.E. Clark and S. T. Love and G. Wu, Phys.Rev.D57, 5912(1998)
- [20] T. E. Clark, M. Nitta and T. ter Veldhuis, Phys.Rev.D67, 085026(2003)
- [21] T. E. Clark, M. Nitta and T. ter Veldhuis, Phys.Rev.D70, 125011(2004)
- [22] E. A. Ivanov and V. I. Ogievetsky, Teor. Mat. Fiz. 25, 164(1975)
- [23] Lu-Xin Liu, Eur.Phys.J.C62:615-623,2009; arXiv:0802.1299 [hep-th]
- [24] P. West, JHEP 0002, 24(2000)
- [25] J. Hughes and J. Polchinski, Nucl. Phys. B278, 147(1986); C. P. Burgess, E. Filotas, M. Klein and F. Quevedo, JHEP 0310, 41(2003); J. Bagger, A. Galperin, Phys. Lett. B336, 25(1994); Phys. Lett. B412, 296(1997); Phys. Rev. D55, 1091(1997)
- [26] S. Bellucci, E. Ivanov and S. Krivonos, Phys. Lett. B460, 348(1999); M. Rocek and A.A.Tseytlin, Phys. Rev. D59, 106001(1999); S. Bellucci, E. Ivanov and S. Krivonos, Fortsch.Phys.48, 19(2000); F. Gonzalez-Rey, L. Y. Park, and M. Rocek, Nucl. Phys.B544, 243(1999)
- [27] I. N. McArthur, hep-th/9908045; S. Bellucci, E. Ivanov, S. Krivonos, Phys.Lett.B482, 233(2000); S. Bellucci, E. Ivanov and S. Krivonos, Nucl. Phys. Proc. Suppl.102, 26(2001); E. Ivanov, Theor. Math. Phys. 129, 1543(2001)
- [28] O. Barwald and P. West, Phys. Lett. B476, 157(2000)
- [29] V.A.Rubakov and M.E.Shaposhnikov, Phys.Lett. B125, (1983)136; E.Papantonopoulos, hep-th/0202044; H.A.Chamblin and H.S.Reall, Nucl.Phys.B562, 133(1999)
- [30] G.Dvali and M.Shifman, Nucl.Phys.B504, 127(1997)

[**31**] N.Maru, N.Sakai, Y.Sakamura and R.Sugisaka, Nucl.Phys.B616,47(2001); Phys.Lett.B496, 98(2000); M.Eto, N.Maru and N.Sakai, Nucl.Phys.B696, 3(2004)